

Identifying research waste from surgical research: a protocol for assessing compliance with the IDEAL framework and recommendations

Jiajie Yu ¹, Fei Shan,² Allison Hirst,³ Peter McCulloch ³, Youping Li,¹ Xin Sun¹

To cite: Yu J, Shan F, Hirst A, *et al*. Identifying research waste from surgical research: a protocol for assessing compliance with the IDEAL framework and recommendations. *BMJ Surg Interv Health Technologies* 2021;**3**:e000050. doi:10.1136/bmjst-2020-000050

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjst-2020-000050>).

JY and FS contributed equally.

Received 04 June 2020
Revised 09 September 2020
Accepted 19 October 2020



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹IDEAL China centre, West China Medical Centre, Chengdu, China
²Gastrointestinal Cancer Center, Peking University Cancer Hospital & Institute, Beijing, China

³Nuffield Department of Surgical Science, University of Oxford, Oxford, UK

Correspondence to

Dr Peter McCulloch;
peter.mcculloch@nds.ox.ac.uk

ABSTRACT

Introduction Approximately £1130 billion was invested in research worldwide in 2016, and 9.6% of this was on biomedical research. However, about 85% of biomedical research investment is wasted. The Lancet published a series to identify five categories relating to research waste and in 2014. Some categories of research waste in surgery are avoidable by complying with the Idea, Development, Exploration, Assessment, Long-term follow-up (IDEAL) framework for it enables researchers to design, conduct and report surgical studies robustly and transparently. This review aims to examine the extent to which surgical studies adhered to the IDEAL framework and estimate the amount of overall research waste that could be avoided if compliance was improved.

Methods We will search for potential studies published in English and between 1 January 2018 and 31 December 2018 via PubMed. Teams of paired reviewers will screen titles, abstracts and full texts independently. Two researchers will extract data from each paper. Data will be collected about general information and specialised information in each stage, and our IDEAL Compliance Appraisal tool will be used to analyse included studies. Descriptive statistics and χ^2 or Fisher's exact tests for comparisons will be presented.

Discussion Our study will provide important information about whether compliance with the specific IDEAL Recommendations has reduced research waste in surgical and therapeutic device studies. And we will identify particular key aspects that are worse and need to focus on improving those in future education.

BACKGROUND

Surgical research is an essential part of clinical research and associated with a number of methodological and practical challenges. The Idea, Development, Exploration, Assessment and Long-term follow-up (IDEAL) Framework and Recommendations were established in 2009, and specify several unique features of clinical studies of surgical interventions, especially for earlier stages of surgical research.¹ The IDEAL pathway integrates proposals regarding the design, conduct and reporting of clinical studies of surgery and has been

widely accepted in studies on surgical intervention, medical devices and physical therapy.²⁻⁴

According to the report prepared by Elsevier for UK's Department for Business, Energy & Industrial Strategy, approximately £1130 billion was invested in research worldwide in 2016, and 9.6% of this was on biomedical research.⁵ However, it is estimated that about 85% of biomedical research investment is wasted.⁶ In 2014, The Lancet published a series to discuss issues relating to research waste.⁷⁻¹¹ These articles identified five categories of research waste. First, research priorities are not based on the needs of potential users (ie, health professionals, patients, policy-makers). Second, weakness is common in the design, conduct and analysis of health research studies (eg, sample size, randomisation, blinding). Third, full information about studies is not accessible (ie, protocol, analyses plan, full study report and participant level data). Fourth, the regulation and management of the conduct of research are often inefficient. Finally, research waste also arises from inadequate reporting of research studies.¹²

Some categories of research waste in surgery (ie, design and reporting flaws) are avoidable by complying with the IDEAL framework.¹² IDEAL enables researchers and authors to design, conduct and report surgical studies in a robust and transparent way.^{13 14} Studies have explored research waste in IDEAL stage 3 (randomised controlled trial, RCT) and found a considerable burden of research waste in surgical RCTs.^{15 16} RCTs, however, only make up a small proportion of published surgical research.¹⁷ Since the IDEAL recommendations cover studies at all stages of the life cycle, and the majority of published studies of safety, efficacy and effectiveness are not randomised trials, better




Figure 1 Flow chart for screening and data extraction. IDEAL, Idea, Development, Exploration, Assessment, Long term.

adherence to IDEAL recommendations could reduce the total sum of research waste in surgery very significantly. We, therefore, decided to conduct a literature survey to examine the extent to which surgical studies adhered to the IDEAL framework to ensure appropriate, safe and efficient translation of new treatments into surgical practice. This will allow estimation of the amount of overall research waste which could be avoided if compliance was improved.

This study aims to assess the avoidable research waste relating to the design, conduct and reporting of studies evaluating innovation, effectiveness, efficacy or harms of surgical operations or therapeutic devices in recently year. Since the research waste associated with methodological and reporting defects in IDEAL stage 3 (RCT) has already been estimated,¹⁶ we will include papers that report studies in stage 1 (idea), stage 2a (development), stage 2b (exploration) and stage 4 (long-term follow-up) according to IDEAL framework.

METHODS

Eligibility criteria

Studies will be included in the review if they meet the following inclusion criteria¹: **participants**: any patient receiving surgical operations or therapeutic devices²; **intervention**: at least one patient/group receiving any kind of surgical operations or therapeutic devices³; **comparison**: alternative surgical approaches and/or non-surgical treatments; no comparator group in some early phase studies⁴; **outcomes**: technical and/or physiological and/or clinical measures of intended effects and/or safety outcomes (including patient-reported outcomes)⁵; **study designs**: non-RCT, prospective cohort study, retrospective cohort study, case-control study, before and after study/interrupted time series, case series and case report.

A surgical operation is defined as the procedure where access is gained via an incision, natural orifice

or percutaneous puncture or one that involves a device used inside the body. Radiological procedures performed without additional surgical intervention are not considered to be within the definition of a surgical study.¹⁶ Procedures testing injection, acupuncture or diagnostic interventions are not eligible, and those that tested effects of medical therapies in patients receiving surgical procedures, such as antibiotics or adjuvant treatments in cancer, will also not be considered. A therapeutic device is defined as a medical device used on humans with therapeutic benefits generally caused through a physical or mechanical effect on the body.

Literature search

We will search for potential publications via PubMed and the search strategy will be developed with the assistance of a specialist librarian. Search terms for surgical intervention and medical therapeutic devices will use those in our published study,¹⁷ and study designs will refer to search filters in Cochrane and the BMJ (online supplemental appendix 1). We will further limit our search to literature published in English and between 1 January 2018 and 31 December 2018. This will allow us to study a very large population of recent surgical studies, and the interval chosen will allow sufficient time for MeSH indexing to be completed before the search for published manuscripts. Citations will be collected by Endnote reference management software.

Screening and selection procedure

An inclusion/exclusion form will be used to screen titles and abstracts. Teams of paired reviewers, trained in the IDEAL framework, will use pilot-tested, standardised forms to screen titles and abstracts independently. We will conduct a pilot study before formal screening and data extraction, during which we will iteratively adjust the protocol and definitions used by reviewers to improve consistency between reviewers, and regular sampling with analysis of the interclass correlation coefficient (ICC) will be used to assess the progress of this process. Any conflicts not resolved by discussion between these reviewers will be referred to the study team for discussion. Full-text versions of papers will be obtained for inclusion in the study. The flow chart for retrieval, screening and exclusions are shown in figure 1.

Data extraction

Data will be extracted independently by at least two assessors for each study. We will randomly select 10% of included papers and evaluate the ICC between reviewers for these papers for both data extraction results and IDEAL compliance tool results. The following information will be collected for each eligible study:

General information

1. Author, date of publication, country of study (correspondent author), journal of publication; number of arms; recruitment setting (institute, national or international); number of centres; sample size.

2. Intervention type (surgical operation or therapeutic devices).
3. Trial design (non-RCT, prospective cohort study, retrospective cohort study, case-control study, before and after study/interrupted time series, cross-sectional study, case series, case report).
4. The study stage (stage 1, stage 2a, stage 2b and stage 4). This will be determined in term of the purpose/research questions of the study with the updated IDEAL Framework and recommendations. The key research questions (purpose) in each stage are: what is the new treatment concept and why is it needed (proof of concept/stage 1)? Has the intervention reached a state of stability sufficient to allow replication by others (development of procedure/stage 2a)? Have the questions that might compromise the chance of conducting a successful RCT been addressed (achieving consensus between surgeons and centres/stage 2b)? Are there any long-term or rare adverse effects or changes in indications or delivery quality over time (surveillance/stage 4).
5. Details of the protocol (reference, Digital Object Identifier (DOI), registration number, supplement).
6. Whether informed consent was obtained or not.

7. Funding source and/or conflict of interest.
8. Whether the authors state they were trying to use the IDEAL recommendations in their study.

Specialised information in each stage

The study will use our IDEAL Compliance Appraisal tool to analyse included studies. Details of the development and validation of the tool will be published separately, and the results of this study will provide further evidence to confirm its validity and reliability. Every domain in each stage is assessed as ‘yes/green’ for a positive result, ‘no/red’ for a negative result or ‘partial yes/yellow’ where neither of these appears appropriate. The details of the appraisal tool see in [table 1](#). We will include a brief narrative report on observed tendencies and trends within the data, including analysis of the areas of best and worst compliance overall, and a comment on any notable differences between the surgery and device subgroups.

Sample size

Two hundred studies (100 on surgical operations and 100 on therapeutic devices) will be included in this study. Studies will be selected at random from the pool

Table 1 IDEAL Compliance Appraisal tool		
Stage	Domains	Option
Stage 1	Idea	
Item 1	Was the patient selection process fully described.	<input type="checkbox"/> Yes <input type="checkbox"/> Partial Yes <input type="checkbox"/> No
Item 2	Did the study fully describe the procedure of interest.	<input type="checkbox"/> Yes <input type="checkbox"/> Partial Yes <input type="checkbox"/> No
Item 3	Did the study clearly state the rationale for the innovation.	<input type="checkbox"/> Yes <input type="checkbox"/> Partial Yes <input type="checkbox"/> No
Item 4	Did the study appropriately report relevant outcome measures.	<input type="checkbox"/> Yes <input type="checkbox"/> Partial Yes <input type="checkbox"/> No
Stage 2a	Development	
Item 1	Did the study fully describe the patient selection process as used in the first case in the 2a study.	<input type="checkbox"/> Yes <input type="checkbox"/> Partial Yes <input type="checkbox"/> No
Item 2	Did the study describe the procedure of interest as it was performed in the first case in the 2a study.	<input type="checkbox"/> Yes <input type="checkbox"/> Partial Yes <input type="checkbox"/> No
Item 3	Did the study report all changes in the design, use of the device or the way of performing the procedure.	<input type="checkbox"/> Yes <input type="checkbox"/> Partial Yes <input type="checkbox"/> No
Item 4	Did the study report key outcome measures sequentially (case by case) to allow evaluation of change.	<input type="checkbox"/> Yes <input type="checkbox"/> No
Item 5	Did the study state if the innovation stable enough to move to stage 2b.	<input type="checkbox"/> Yes <input type="checkbox"/> Partial Yes <input type="checkbox"/> No
Stage 2b	Assessment	
Item 1	Was there a unified description of the intervention and accepted variation.	<input type="checkbox"/> Yes <input type="checkbox"/> Partial Yes <input type="checkbox"/> No
Item 2	Did the study define the measures used to evaluate the quality of the procedure or device use.	<input type="checkbox"/> Yes <input type="checkbox"/> No
Item 3	Did the study provide information about the learning curve assessment.	<input type="checkbox"/> Yes <input type="checkbox"/> Partial Yes <input type="checkbox"/> No
Item 4	Did the study provide a qualitative evaluation of the stakeholder values.	<input type="checkbox"/> Yes <input type="checkbox"/> Partial Yes <input type="checkbox"/> No
Item 5	Did the study describe a pre-planned review of results and discussion to achieve the agreement on the stage 3 trial.	<input type="checkbox"/> Yes <input type="checkbox"/> Partial Yes <input type="checkbox"/> No
Item 6	Did the study report the use of the primary outcome effect estimate to perform power calculations for possible RCTs.	<input type="checkbox"/> Yes <input type="checkbox"/> No
Stage 4	Long-term follow-up.	
Item 1	Did the study describe the data source.	<input type="checkbox"/> Yes <input type="checkbox"/> No
Item 2	Did the study clearly and precisely describe the definition of the dataset.	<input type="checkbox"/> Yes <input type="checkbox"/> Partial Yes <input type="checkbox"/> No
Item 3	Did the study conduct appropriate analysis of data.	<input type="checkbox"/> Yes <input type="checkbox"/> Partial Yes <input type="checkbox"/> No
Item 4	Did the study report the missing data of the dataset.	<input type="checkbox"/> Yes <input type="checkbox"/> Partial Yes <input type="checkbox"/> No

RCT, randomised controlled trial.



of search literature for screening and data extraction. A sample of studies will be selected randomly in a 1:1 ratio between studies of operations and therapeutic devices. If the number of eligible studies is insufficient after screening and data extraction, sampling of the identified studies will continue, until 100 of each type is selected.

Statistical analysis

We will estimate the avoidable research waste in each sample by comparing the actual number of 'green' item assessments versus the total number of items. We will estimate the potential contribution to reducing overall research waste if IDEAL were perfectly complied with by considering the percentage of all research registered on PubMed represented by the outcome studies selected by our search, and the proportion of this represented by different subgroups (eg, device vs procedure studies; prospective vs retrospective design).

For all descriptive analyses, we will use frequencies (and percentages) for dichotomous variables, and mean (and SD) or median (and range) or median (and first quartile, third quartile) for continuous variables. Either χ^2 or Fisher's exact tests will be used for the comparison between studies in surgical intervention and medical therapeutic devices. All analyses will be performed by IBM SPSS Statistics V.26.

DISCUSSION

The Lancet published one of its most provocative Commentaries in 1996 and labelled 'surgical research' as 'comic opera'.¹⁸ This critique was based largely on surgical studies in the early stages. Our study will provide important information about whether compliance with the specific IDEAL Recommendations has reduced research waste in surgical and therapeutic device studies, especially those studies in earlier stages of surgical research. Additionally, we will identify particular aspects of study design, conduct and reporting that need improvement, usefully directing the focus of future education.

To our knowledge, this study is the first study assessing research waste in non-randomised studies and will include all specialties and all stage of the life cycle (from innovation to long-term follow-up), which should lead to better generalisability of the findings.

There are some limitations in this study. First, of the five key elements of research waste in the Lancet series, we are not able to assess waste due to failure to identify the research priority, research management issues and lack of fully accessible research, because published reports contain no information about these problems. Our methodological study, therefore, focuses on avoidable design, conduct and reporting errors. Second, our study is based on published information only, and may not reflect the actual implementation of surgical research.

Twitter Allison Hirst @IDEALCollab

Contributors JY, FS and PM conceived and initiated the study and designed the protocol. JY and FS wrote the first draft of this manuscript. JY, FS, AH, YL, XS and PM critically revised the protocol and manuscript.

Funding The IDEAL Collaboration is supported by the Oxford NIHR Biomedical Research Centre, an unrestricted Educational Grant from Medtronic as well as other educational grants and meetings income. JY was supported by a grant from 'National Natural Science Foundation of China' (project No: 71704120), 'National Science and Technology Major Projects' (project No: 2018Z×09201009–005–003) and International visiting programme for Excellent Young Scholars of SCU.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Jiajie Yu <http://orcid.org/0000-0003-4319-0820>

Peter McCulloch <http://orcid.org/0000-0002-3210-8273>

REFERENCES

- McCulloch P, Altman DG, Campbell WB, *et al*. No surgical innovation without evaluation: the ideal recommendations. *Lancet* 2009;374:1105–12.
- Hirst A, Philippou Y, Blazeby J, *et al*. No surgical innovation without evaluation: evolution and further development of the ideal framework and recommendations. *Ann Surg* 2019;269:211–20.
- Sedrakyan A, Campbell B, Merino JG, *et al*. IDEAL-D: a rational framework for evaluating and regulating the use of medical devices. *BMJ* 2016;353:i2372.
- Beard D, Hamilton D, Davies L, *et al*. Evidence-based evaluation of practice and innovation in physical therapy using the IDEAL-Physio framework. *Phys Ther* 2018;98:108–21.
- Department for Business EIS. *International comparative performance of the UK research base 2016*, 2016.
- Chalmers I, Glasziou P. Avoidable waste in the production and reporting of research evidence. *Lancet* 2009;374:86–9.
- Al-Shahi Salman R, Beller E, Kagan J, *et al*. Increasing value and reducing waste in biomedical research regulation and management. *Lancet* 2014;383:176–85.
- Chalmers I, Bracken MB, Djulbegovic B, *et al*. How to increase value and reduce waste when research priorities are set. *Lancet* 2014;383:156–65.
- Chan A-W, Song F, Vickers A, *et al*. Increasing value and reducing waste: addressing inaccessible research. *Lancet* 2014;383:257–66.
- Ioannidis JPA, Greenland S, Hlatky MA, *et al*. Increasing value and reducing waste in research design, conduct, and analysis. *Lancet* 2014;383:166–75.
- Glasziou P, Altman DG, Bossuyt P, *et al*. Reducing waste from incomplete or unusable reports of biomedical research. *Lancet* 2014;383:267–76.
- Macleod MR, Michie S, Roberts I, *et al*. Biomedical research: increasing value, reducing waste. *Lancet* 2014;383:101–4.
- Khachane A, Philippou Y, Hirst A, *et al*. Appraising the uptake and use of the ideal framework and recommendations: a review of the literature. *Int J Surg* 2018;57:84–90.
- Main BG, Blencowe NS, Howes N, *et al*. Protocol for the systematic review of the reporting of transoral robotic surgery. *BMJ Open* 2018;8:e019198.
- Chapman SJ, Shelton B, Mahmood H, *et al*. Discontinuation and non-publication of surgical randomised controlled trials: observational study. *BMJ* 2014;349:g6870.
- Chapman SJ, Aldaffaa M, Downey CL, *et al*. Research waste in surgical randomized controlled trials. *Br J Surg* 2019;106:1464–71.
- McCulloch P, Feinberg J, Philippou Y, *et al*. Progress in clinical research in surgery and ideal. *Lancet* 2018;392:88–94.
- Horton R. Surgical research or comic opera: questions, but few answers. *Lancet* 1996;347:984–5.

Appendix 1 Search strategy

Step	Query	Description
#1	Search (surgery[MeSH Terms]) OR surgical procedures, operative[MeSH Terms]	MeSH search for "SURGICAL OPERATIONS"
#2	Search ((((((((((Surgical[Title/Abstract]) OR Operation[Title/Abstract]) OR operative[Title/Abstract]) OR operative therapy[Title/Abstract]) OR operative procedure[Title/Abstract]) OR invasive procedure[Title/Abstract]) OR peroperative procedure[Title/Abstract]) OR perioperative procedure[Title/Abstract]) OR preoperative procedure[Title/Abstract]) OR intraoperative procedure[Title/Abstract]) OR surgery[Title/Abstract]) OR procedure[Title/Abstract]) OR Surgical Procedure[Title/Abstract]	Keyword search for "SURGICAL OPERATIONS"
#3	Search (equipment and supplies[MeSH Terms])	MeSH search for "THERAPEUTIC DEVICES"
#4	Search ((((((equipment[Title/Abstract]) OR supplies[Title/Abstract]) OR apparatus[Title/Abstract]) OR instrument[Title/Abstract]) OR inventor*[Title/Abstract]) OR Medical Device[Title/Abstract]) OR Device[Title/Abstract]	Keyword search for "THERAPEUTIC DEVICES"
#5	Search drug therapy[MeSH Terms]	MeSH search for "DRUGS"
#6	Search (((Drug Therap*[Title/Abstract]) OR Drug treatment[Title/Abstract]) OR Chemotherap*[Title/Abstract]) OR Pharmacotherap*[Title/Abstract]) OR pharmacologic[Title/Abstract]	Keyword search for "DRUGS"
#7	Search #1 OR #2	
#8	Search #3 OR #4	
#9	Search (#5) OR #6	
#10	Search #7 NOT #9	
#11	Search #8 NOT #9	
#12	"case control studies"[Mesh] OR "Cohort Studies"[Mesh] OR "Longitudinal Studies"[Mesh] OR "retrospective studies"[Mesh] OR "Follow-Up Studies"[Mesh] OR "prospective studies" [Mesh] OR Cohort[tiab] OR case control[tiab] OR longitudinal[tiab] OR follow up[tiab] OR prospective*[tiab] OR retrospective*[tiab] OR nonrandom*[tiab] OR comparison group*[tiab] OR control group*[tiab] OR "Registries"[Mesh] OR registry[tiab] OR registries[tiab]	search for "NON-RANDOMIZED STUDY"
#13	Search (((("Case Reports" [Publication Type]) OR "Single-Case Studies as Topic"[Mesh])) OR ("case	search for "CASE STUDY"

	report"[Title/Abstract] OR "case reports"[Title/Abstract])) OR "case series"[Title/Abstract] OR "case study") OR "case studies"[Title/Abstract]	
#14	(systematic review [ti] OR meta-analysis [pt] OR meta-analysis [ti] OR systematic literature review [ti] OR systematic review [tw] OR (systematic review [tiab] AND review [pt]) OR meta synthesis [ti] OR integrative review [tw] OR integrative research review [tw] OR rapid review [tw] OR consensus development conference [pt] OR practice guideline [pt] OR cochrane database syst rev [ta] OR acp journal club [ta] OR health technol assess [ta] OR evid rep technol assess summ [ta] OR drug class reviews [ti]) OR (clinical guideline [tw] AND management [tw]) OR meta analy*[ti] OR metaanaly*[ti] OR meta analysis[pt]	Keyword search for "SYSTEMATIC REVIEW"
#15	"Meta-Analysis as Topic"[Mesh] OR "Review Literature as topic"[Mesh]	MeSH search for "SYSTEMATIC REVIEW"
#16	Search (((((((Clinical Conference [Publication Type] OR Comment [Publication Type] OR Congress [Publication Type] OR Editorial [Publication Type] OR Letter [Publication Type] OR Clinical Trial Protocol [Publication Type] OR Clinical Trial, Veterinary[Publication Type])) OR ((Conference[Title/Abstract] OR conferences[Title/Abstract]))) OR ((Comment[Title/Abstract] OR comments[Title/Abstract]))) OR Congress[Title/Abstract] OR Consensus[Title/Abstract] OR ((Editorial[Title/Abstract] OR editorials[Title/Abstract] OR Letter[Title/Abstract] OR Protocol[Title/Abstract] OR protocols[Title/Abstract]))) OR Clinical Trial, Veterinary[Publication Type]	Search "EDITORIAL"
#17	Search #12 OR #13	
#18	Search #14 OR #15 OR #16	
#19	Search #10 AND #17 NOT #18	Surgical operation
#20	Search #11 AND #17 NOT #18	Therapeutic devices
#21	Search English[Language]	
#22	Search animals [Mesh]	
#23	Search humans [Mesh]	
#24	Search #22 NOT #23	
#25	Search ("2018/1/1"[Date - Publication] : "2018/12/31"[Date - Publication])	
#26	Search #19 AND #21 NOT #24	
#27	Search #25 AND #26	Surgical operation
#28	Search #20 AND #21 NOT #24	
#29	Search #28 AND #26	Therapeutic devices

